

Geodetic Graphs of Diameter Two and Some Related Structures*

RAFFAELE SCAPELLATO

*Dipartimento di Matematica, Università di Parma,
43100 Parma, Italy*

Communicated by the Managing Editors

Received March 30, 1985

We show a connection between 2-connected geodetic graphs of diameter two and certain geometric structures; we apply this result, proving that these graphs fall into three classes: (1) strongly regular graphs with $\mu = 1$; (2) pyramids; (3) π -graphs. (The definitions for (1), (2), and (3) are given in the text.) We also give some results on graphs of type (3), including some non-existence conditions. Finally, we consider a more general class of graphs, showing a connection between some of them and Sperner spaces. © 1986 Academic Press, Inc.

INTRODUCTION

A (finite, simple, undirected) graph is said to be *geodetic* if every pair of its vertices is connected by one and only one shortest path. Recently many authors have worked on geodetic graphs (see, e.g., [2, 7, 10, 11, 12]).

In the first section of this paper we give constructions and results on some classes of graphs, which we will apply in what follows. Next we present a classification of geodetic 2-connected graphs of diameter two. We also show (section 2) how geodetic graphs of diameter two with certain parameters are obtained from affine planes.

Finally, in Section 3 we deal with a larger class of graphs (*F*-geodetic), which are a natural generalization of both geodetic and distance-regular graphs; among other things, it is proved that there is a connection between some of these graphs and Sperner spaces.

1. GEODETIC GRAPHS OF DIAMETER TWO

1. In this paper we follow, when it is not otherwise specified, the terminology of [9]; we also suppose all sets in this paper to be *finite*.

* Research supported by the Italian MPI.

If G is a graph, then $V(G)$ ($E(G)$) is the vertex set (the edge set) of G . We denote by $(x; y)$ the unordered pair (edge) which consists of x and y .

A *clique* of a graph G is a maximal complete subgraph of G , with at least three vertices; a *clique-vertex* is a vertex which belongs to some clique.

Let G be a graph, let $x, y \in V(G)$. We denote by $d_G(x, y)$, or $d(x, y)$, the distance between x and y ; moreover, we put

$$S^-(x, y) = \{z \in V(G) \mid (x; z) \in E(G), d(y, z) = d(x, y) - 1\},$$

$$S^+(x, y) = \{z \in V(G) \mid (x; z) \in E(G), d(y, z) = d(x, y) + 1\}.$$

A *distance-regular* graph is a connected graph G such that there are non-negative integers $b_0, b_1, \dots, b_h, c_0, c_1, \dots, c_h$ (with $c_0 = b_h = 0$ and $c_1 = 1$), where h is the diameter of G , such that for $x, y \in V(G)$ with $d(x, y) = j$ it is $|S^-(x, y)| = c_j$, $|S^+(x, y)| = b_j$ (see, e.g., [1]).

A *strongly regular* graph is a regular graph G for which there are non-negative integers λ and μ , such that the number of common neighbours of two different vertices is λ or μ , accordingly to whether they are or are not adjacent. It is known that these are exactly the distance-regular graphs of diameter two.

For a graph G , a shortest path between two vertices is, as in [3, 13], called a *geodesic*.

Let G be a connected graph of diameter h , and let F be a map from $\{1, 2, \dots, h\}$ to the set of positive integers. We say that G is F -geodetic if, for every $i \in \{1, 2, \dots, h\}$ and for every $x, y \in V(G)$, with $d(x, y) = i$, the number of geodesics between x and y is exactly $F(i)$.

Geodetic graphs are, therefore, F -geodetic graphs for which the map F is the constant 1; we call *proper* an F -geodetic but not a geodetic graph.

Let H be a graph of diameter $h \neq 1$, such that, for every $x \in V(H)$, there is a $y \in V(H)$ with $d(x, y) = h$. If there is a surjection $\varphi: V(H) \rightarrow A$ (where A is a set, disjoint from $V(H)$), such that $\varphi(x) = \varphi(y)$ if and only if $d(x, y) = h$ or 0, then H is said, with [4], to be *antipodal*. The map φ is said to be a *natural coloring* of H .

We call *fully antipodal* an antipodal graph H such that for $x, y \in V(H)$ with $\varphi(x) = \varphi(y)$ there is a $y' \in V(H)$ with $\varphi(y) = \varphi(y')$ and $(x, y') \in E(H)$. We call *perfectly antipodal* a fully antipodal graph H such that for every $x, y \in V(H)$ with $\varphi(x) = \varphi(y)$ there is a $z \in V(H)$ with $(x; z), (z; y) \in E(H)$. Such graphs are characterized by Theorem 2.

For the reader's convenience, we recall below a list of results shown in [13].

In what follows, G is a 2-connected geodetic graph of diameter two.

1. If G is regular, it has no cliques, or all vertices are clique-vertices [13, Theorems 5.3 and 5.6].

2. If G is not regular there are two integers m, n , with $1 < m < n$, such that any vertex of G has degree m or n : clique-vertices and vertices whose distance from each clique is 2 have degree n , and the others have degree m [13, Theorem 5.6]; the number of vertices of G is $mn + 1$ [13, Theorem 5.2]; all cliques of G have $k = n - m + 2$ vertices [13, Theorem 5.5]; two vertices of degree m cannot be adjacent [13, Theorems 5.5 and 5.6].

3. If two clique-vertices are adjacent, they lie in the same clique [13, Lemma 5.7].

From now on we will use these results without mention; moreover, if m, n are integers, with $1 < m < n$, we denote by $\mathcal{G}(m, n)$ the class of all geodetic graphs as in result 2 above.

2. We now introduce some definitions.

Let P and R be disjoint sets, let $I \subseteq P \times R$ be a relation, let $\parallel \subseteq R \times R$ be a reflexive and symmetric relation; we call *points* the elements of P , *lines* the elements of R , *incidence* the relation I , *parallelism* the relation \parallel .

We say that the ordered quadruple (P, R, I, \parallel) is a π -space if the following axioms hold:

- (A1) Two different points are incident with one and only one line.
- (A2) For every point x and every line r , there is a unique line parallel to r and incident with x .
- (A3) If two lines are neither parallel nor incident with the same point, there is a unique line parallel to both.
- (A4) There are at least two points, which are not incident with the same line; every line is parallel to at least two other lines, parallel to each other.

Moreover, we say that (P, R, I, \parallel) is a *Sperner space* (see, e.g., [8]) if it verifies (A1), (A2), and (A4), and there is a nonnegative integer k such that each line is incident with exactly k points, and \parallel is transitive.

If (P, R, I, \parallel) is an affine plane (of order $q \geq 3$), it is of course a π -space; moreover, let D be the set of equivalence classes of \parallel ; put $R' = R \cup D$, $\parallel' = \parallel \cup \{(r, r') | r \in r', r \in R, r' \in D\}$. Then (P, R', I, \parallel') is again a π -space, which we call a *modified affine plane*.

We state the following without proof:

Let G be a 2-connected, non-regular, geodetic graph of diameter 2, without vertices at distance 2 from each clique. Let R be the set of all clique-vertices of G ; put $P = V(G) \setminus R$, $I = \{(x, r) \in P \times R | (x, r) \in E(G)\}$, $\parallel = \{(r, r') \in R \times R | (r, r') \in E(G) \text{ or } r = r'\}$. Then (P, R, I, \parallel) is a π -space.

Conversely, if (P, R, I, \parallel) is a π -space, by $V(G) = P \cup R$ and $E(G) = \{(x; r) | (x, r) \in I\} \cup \{(r; r') | (r, r') \in \parallel \text{ and } r \neq r'\}$, we obtain a 2-connected, non-regular, geodesic graph of diameter 2, without vertices at distance 2 from each clique.

DEFINITION 1. A π -graph is a graph which satisfies the above conditions.

We now introduce the following notation, which is used throughout this paper.

Let $\mathcal{P} = (P, R, I)$ be a projective plane of order $q \geq 3$, and let σ be a polarity of \mathcal{P} . A point x of \mathcal{P} is said to be *absolute* for σ if $\sigma(x) = x$ (see [5]). We denote by $G(\mathcal{P}, \sigma)$ the graph G defined by $V(G) = P$ and $E(G) = \{(x; y) | xI\sigma(y)\}$. Moreover, if the *absolute points* of σ are exactly the points incident with a line r , we denote by $H(\mathcal{P}, \sigma)$ the *section graph* of G defined by the set of non-absolute points different from $\sigma(r)$.

The graph $G(\mathcal{P}, \sigma)$ (the graph $H(\mathcal{P}, \sigma)$) will be called *projective* (restricted projective).

LEMMA 1. Let H be a fully antipodal graph, and let $\varphi: V(H) \rightarrow A$ be a natural coloring of H . Then H is complete multipartite or it is regular of degree $|A| - 1$ and diameter 3.

Proof. Let h be the diameter of H : we show that $h \leq 3$. By contradiction, we suppose $h > 3$.

If $b, b' \in V(H)$, with $d(b, b') = 2$, then $\varphi(b) \neq \varphi(b')$, so by definition there is a z with $(z; b') \in E(H)$ and $\varphi(z) = \varphi(b)$. But now z and b have the same color and their distance is less than the diameter, contradicting the definition of antipodal graph. Then $h \leq 3$ and, by definition again, $h \neq 1$.

If $h = 2$, by [4, Corollary 3.6] the graph H is complete multipartite.

Consider now the remaining case $h = 3$. Since each vertex b of H is adjacent to at least one vertex of every color different from $\varphi(b)$, the degree of b is at least $|A| - 1$; since two vertices of the same color must have distance 3, the degree of b is exactly $|A| - 1$. The lemma is now proved.

EXAMPLE. The graph illustrated by Fig. 1 (colored with 1, 2, 3, 4) is fully antipodal of degree 3 and diameter 3, and also F -geodesic (with $F(2) = 1$, $F(3) = 2$). It is not distance-regular, and of course not perfectly antipodal.

THEOREM 2. A graph is perfectly antipodal if and only if it is a restricted projective graph, or a complete multipartite graph with at least three parts.

Proof. By Lemma 1, we must only show that a graph of diameter 3 is perfectly antipodal if and only if it is a restricted projective graph.




FIGURE 1

Let H be a perfectly antipodal graph of diameter 3, $\varphi: V(H) \rightarrow A$ be a natural coloring of H : then H has diameter 3 and it is regular of degree $q = |A| - 1$ (Lemma 1).

Let $c \notin A \cup V(H)$, $P = V(H) \cup A \cup \{c\}$. For $x \in P$ we state:

$$\sigma(x) = \begin{cases} \{y \in V(H) | (x, y) \in E(H)\} \cup \{\varphi(x)\} & \text{if } x \in V(H) \\ \{y \in V(H) | \varphi(y) = x\} \cup \{x, c\} & \text{if } x \in A \\ A & \text{if } x = c. \end{cases}$$

Finally, we put $R = \{\sigma(x) | x \in P\}$. Now we consider the incidence structure $\mathcal{P} = (P, R, \in)$. It is easy to show that two points are incident with at least one line.

We show that every line of \mathcal{P} is incident with exactly $q + 1$ points. First, it is clear that it is true for all lines $\sigma(x)$, with $x \in V(H) \cup \{c\}$.

For $a \in A$, consider the line $\sigma(a)$. Let b be a vertex of H : every line for b must intersect $\sigma(a)$. Let $\sigma(x)$, $\sigma(y)$ be two lines incident with b and intersecting $\sigma(a)$ in the same point a' . Then $\varphi(x) = \varphi(y) = a'$; since b is adjacent to both x and y , we have $d(x, y) \neq 3$. The hypotheses on H thus imply $x = y$. Then every line by b intersects $\sigma(a)$ in a unique point, so the points of $\sigma(a)$ are in one-to-one correspondence with the lines for b , thus are $q + 1$.

It is now easy to see that \mathcal{P} is a projective plane of order q and that σ is a polarity of \mathcal{P} , whose absolute points are the elements of A , and $H = H(\mathcal{P}, \sigma)$.

Conversely, let $H = H(\mathcal{P}, \sigma)$ be a restricted projective graph, let A be the set of all absolute points of σ , let c be the pole of the line incident with all the points of A , and let $\varphi: V(H) \rightarrow A$ be the map which maps $b \in V(H)$ in the intersection of A with the line for b and c .

We show that H has diameter 3 and is antipodal. Let $b, b' \in V(H)$, with $b \neq b'$, such that b, b', c lie in the same line (note that in the definitions of projective and restricted projective graph we suppose that $q \geq 3$: so in a line for c there are at least two points not in A). Since their polars must intersect in a point of the polar of c , i.e., of A , in H no vertex is adjacent to both: so $d(b, b') \geq 3$.

With similar arguments it is now easy to conclude the proof.

DEFINITION II. A graph G is a pyramid of type h if there is a graph H of diameter h and there is a map φ from $V(H)$ to a set A (disjoint from $V(H)$), such that:

- (1) the graph H is perfectly antipodal and φ is a natural coloring of H , or H is a complete graph and φ is a bijection;
- (2) there is a $c \in V(G)$ such that $V(G) = V(H) \cup A \cup \{c\}$ and $E(G) = E(H) \cup \{(b; \varphi(b)) | b \in V(H)\} \cup \{(a; c) | a \in A\}$.

THEOREM 3. Let G be a 2-connected geodetic graph of diameter 2. If c is a vertex of G , whose distance from all cliques is two, then G is a pyramid of type 1 or 3.

Proof. Let $A = \{x \in V(G) | (x; c) \in E(G)\}$, $B = V(G) \setminus (A \cup \{c\})$; let H be the section graph of G defined by B .

Since the elements of B are at distance 2 from c , for every $b \in B$ there is an $a \in A$ adjacent to b , which must be unique (because if it is not so, c and b are in a circuit of length 4, thus are adjacent). Now we put $\varphi(b) = a$, obtaining a map from B to A .

If H is a complete graph, then the map φ is injective: thus, for $b \in B$, $\varphi(b)$ has degree 2, so [13, Theorem 5.7 and Corollary 4] G is a pyramid of Type 1.

If H is not complete, let $b, b' \in B$ with $b \neq b'$ and $\varphi(b) = \varphi(b')$. If $(b; b') \in E(H)$, we have $(b; b') \in E(G)$, so b, b' and $\varphi(b)$ are in the same clique; the same holds if $d_H(b, b') = 2$. But this is a contradiction, since $\varphi(b)$ is not a clique-vertex: thus $d_H(b, b')$ is greater than 2.

Let x be an element of B , adjacent to b ; the vertex y such that (x, y, b') is a geodesic cannot be $\varphi(b)$, so $y \in B$. This proves that $d_H(b, b') = 3$.

Conversely, if $b, b' \in B$ have distance 3 in H , from the fact that G has diameter 2 it follows that $\varphi(b) = \varphi(b')$. Thus H is antipodal and φ is a natural coloring of it.

It is now easy to conclude the proof.

COROLLARY 4. *The 2-connected geodetic graphs of diameter two fall in the following classes:*

- (1) *strongly regular graphs with $\mu = 1$;*
- (2) *pyramids of type 1 or 3;*
- (3) *π -graphs.*

2. ON π -GRAPHS

3. We call *principal* a line of a π -space which intersects every line which is not parallel to it. This definition is analogous to that given in [8] for weak affine spaces.

LEMMA 5. *Let S be a π -space, without lines incident with no points, and let G be the π -graph associated to S .*

If $G \in \mathcal{G}(m, n)$, the following hold:

- (1) *a line is principal iff it has exactly $m - 1$ points;*
- (2) *if a line is principal, all lines parallel to it are also principal;*
- (3) *if there are two principal lines which are not parallel, then S is an affine plane.*

Proof. If the line r has $m - 1$ points, let s be a line which does not intersect r . Counting the lines by a point x of s it is clear that s is parallel to r , and r is principal; the other part of (1) is trivial. Thus (1) follows.

Let r be a principal line. By (1) the number of lines parallel to r is $n - m + 1 = k - 1$. Therefore r is in a unique clique.

We suppose, by contradiction, that $s \parallel r$ is *not* principal; let s' be a line of S , such that s and s' are neither parallel nor incident with the same point of S . Let s'' be the parallel to s and s' .

Since r is principal it follows that $s'' \parallel r$, and similarly $s' \parallel r$: then $s \parallel s'$, contradicting the hypothesis on s' . Thus (2) follows.

Now (3) follows easily from (1) and (2).

The next theorem has been proved, in a different way, in [6, Lemma 3.6].

THEOREM 6. *Let $q \geq 4$ be an integer. A graph G is in $\mathcal{G}(q + 1, 2q - 1)$ if and only if G is a π -graph, whose π -space is an affine plane of order q .*

Before the proof, we remark that if m, n are integers, with $2 < m < n - 1$, the class $\mathcal{G}(m, n)$ consists only of π -graphs; on the other hand in pyramids of type 1 it is $m = 2$, and in pyramids of type 3 it is $m = n - 1$ (since such a pyramid is a projective graph; see Theorem 2).

Proof. Let $G \in \mathcal{G}(q+1, 2q-1)$. Since q is greater than 3, the above remark shows that G is a π -graph. Let $S = (P, R, I, \parallel)$ be the associate π -space.

If two elements of R are adjacent in G , then they lie in the same clique, so the number of elements of R adjacent to a given $r \in R$ is a multiple of $k-1$ (where k is the number of the vertices of a clique).

In our case it is $m = q+1$, $n = 2q-1$, so $k-1 = q-1$. Thus r has $q-1$ or $2(q-1)$ neighbours in R , i.e., respectively, it has q or one neighbours in P .

Let x, y , and z be three different points not in a line. The lines by x and y and by x and z have at least two points, then they have q points: because $q = m-1$, S is an affine plane (Lemma 5).

Since the converse is straightforward, the assertion is proved.

THEOREM 7. *Let $q \geq 4$ be an integer. A graph G is in $\mathcal{G}(q+1, 2q)$ if and only if G is a π -graph, whose π -space is a modified affine plane of order q .*

Proof. Let $G \in \mathcal{G}(q+1, 2q)$. By the hypothesis on q and the remark before the proof of Theorem 6, it is clear that G is a π -graph: let $S = (P, R, I, \parallel)$ be its π -space, let R' be the set of all lines incident with some point, let $H = R \setminus R'$, and let \parallel' be the restriction of \parallel to R' ; we prove that $S' = (P, R', I, \parallel')$ is an affine plane.

Given $r \in R$, since the degree of r in G is $2q$ and since r is in at least one clique, from $k = 2q - (q+1) + 2 = q+1$ (where k is the number of elements of a clique) it follows that r is adjacent to q element of P or to none: so r is incident to no points or to exactly q points. Then every line of S' has q points. Reasoning as in Theorem 6 we conclude that S' is an affine plane.

Since the converse is straightforward, the assertion follows.

4. The graphs considered in Theorems 6 and 7 do not exhaust the class of π -graphs: other examples are the projective graphs $G(\mathcal{P}, \sigma)$, where the absolute points of σ are not collinear.

LEMMA 8. *Let S be a π -space, whose π -graph is in $\mathcal{G}(m, n)$. If in S there are principal lines, one of the following holds:*

- (1) *the π -space S is an affine plane of order $m-1$;*
- (2) *each non-principal line of S is incident with exactly $k = n - m + 2$ points.*

Proof. If S is an affine plane of order q , we have $q = m-1$ (Theorem 6), so (1) holds.

Now we suppose that S is not an affine plane. Let r be a principal line of S ; by Lemma 5 there are exactly k lines parallel to r . A non-principal line s

is not parallel to any of these lines, so it intersects each principal line; on the other hand, each point of s lies in some line parallel to r , hence in a principal line. Thus (2) holds and the lemma is proved.

Applying the preceding results, we prove

THEOREM 9. *Let m and n be positive integers, with $m \geq 5$ and $3m \leq 2n + 2$. Then $\mathcal{G}(m, n)$ is non-empty if and only if n is equal to $2(m - 1)$ or $2(m - 1) - 1$ and there is an affine plane of order $m - 1$.*

Proof. The "if" part is an easy consequence of Theorems 6 and 7. Let us suppose that $\mathcal{G}(m, n)$ is non-empty.

Since by definition the elements of $\mathcal{G}(m, n)$ are non-regular graphs, in $\mathcal{G}(m, n)$ there are only pyramids and π -graphs. As observed before the proof of Theorem 9, if in $\mathcal{G}(m, n)$ there are pyramids, it is $(m, n) = (2, n)$ or $(m, n) = (m, m + 1)$. Both cases contradict our hypothesis on n, m (in the second it is $2n + 2 = 2m + 4$, so $m \leq 4$). Therefore, all elements of $\mathcal{G}(m, n)$ are π -graphs.

Let $G \in \mathcal{G}(m, n)$, and let S be the π -space of G . It is easily seen that the points on a line r of S are $n - t(k - 1)$, where t is the number of cliques of G containing r . For $t \geq 3$ we have $n - t(k - 1) \leq n - 3(k - 1) = 3m - 2n - 3 < 0$, so a line of S is in at most two cliques.

A line of S has, therefore, $m - 1$ or $2m - n - 2$ points. By contradiction, we now suppose that no line has $m - 1$ points.

Then S is a BIB-design (with $\lambda = 1$); putting $v = |P|$, we have, therefore, $(v - mn - 1)(2m - n - 2) = vm$, $v - 1 = m(2m - n - 3)$.

Performing some calculations, from the above equalities we have $2(m^2 + n^2) + 5(mn - 1) + 7(n - m) = 0$, a contradiction since n is greater than m . Then, there is a line incident with $m - 1$ points.

If in S no line is incident with no points and if S is not an affine plane, in S there are principal lines (Lemma 5), so $k = 2m - n - 2$ (Lemma 8), so $n - m + 2 = 2m - n - 2$, i.e., $2n + 4 = 3m$, contradicting the hypothesis on m and n . Then if no line is incident with no points, then S is an affine plane and $n = 2(m - 1) - 1$.

Finally, if there is a line incident with no points, then $2m - n - 2 = 0$, i.e., $n = 2(m - 1)$; this proves that there is an affine plane of order $m - 1$ (Theorem 7), completing the proof.

3. ON F -GEODETIC GRAPHS

5. A simple characterization of F -geodetic graphs is given by

THEOREM 10. *A graph G of diameter h is F -geodetic for some F if and only if there are positive integers $c_1 = 1, c_2, \dots, c_h$ such that $\text{diam}(G) = h$ and,*

for every $j \in \{1, 2, \dots, h\}$, if $d(x, y) = j$ then $|S^-(x, y)| = c_j$. Moreover, $F(j) = c_j \cdot F(j-1)$ for each $j \in \{2, 3, \dots, h\}$.

Proof. Suppose that G is F -geodetic; let $j \in \{1, 2, \dots, h\}$. For $x, y \in V(G)$ with $d(x, y) = j$, for every $z \in S^-(x, y)$ there are $F(j-1)$ geodesics between z and y ; adding x to each of them we obtain the same number of geodesics between x and y .

Since every geodesic between x and y must intersect $S^-(x, y)$ in a singleton, the geodesics between x and y are $|S^-(x, y)| \cdot F(j-1)$ in all. Therefore $F(j) = |S^-(x, y)| \cdot F(j-1)$, i.e., $|S^-(x, y)| = F(j)/F(j-1)$. This proves that there are c_1, c_2, \dots, c_h with the requested property.

With a similar argument one can prove that if there are such integers, then G is F -geodetic. The last part of the assertion is now obvious.

PROPOSITION 11. Let $S = (P, R, I, \parallel)$ be a Sperner space, in which each line is incident with k points; suppose that S is an affine plane. Put $V(G) = P \cup R$ and

$$E(G) = \{(r, r') | r, r' \in R, r \neq r', r \parallel r'\} \cup \{(x, r) | x \in P, r \in R, xIr\};$$

then the graph G has diameter 3 and it is F -geodetic, with $F(2) = 1$, $F(3) = 2k$.

Proof. Let $a, b \in V(G)$; it is easy to see that:

—if $a, b \in P$, then $d(a, b) = 2$ and there is a unique geodesic between a and b ;

—if $a \in P, b \in R$ (or $a \in R, b \in P$), then $d(a, b) \leq 2$ and there is a unique geodesic between a and b .

Now we suppose $a, b \in R$. If $a \parallel b$ it is clear that a and b are adjacent in G ; if there is a point x , incident with both a and b , then (a, x, b) is the unique geodesic between them.

There remains the case in which a and b are neither parallel nor incident with the same point: but then $d(a, b) \geq 3$. Moreover, if x is incident to a , let c be the parallel to b by x , then (a, x, c, b) is clearly a geodesic: so $d(a, b) = 3$. For each y incident to b , let d be the parallel to a by y ; then (a, d, y, b) is a geodesic.

It is easily seen that all geodesics between a and b are obtained in this way; since every line has k points, these are $2k$.

As S is not an affine plane, we conclude that G has diameter 3, and this completes the proof.

We call *minimally F -geodetic* an F -geodetic graph G such that, for every $B \subseteq V(G)$, if $B \neq V(G)$ then the section graph of G defined by B is not F -geodetic (with the same map F , hence with the same diameter).

THEOREM 12. *Every proper, minimally F -geodetic graph is 2-connected.*

Proof. Namely let G be a proper, minimally F -geodetic graph of diameter h ; we suppose, by contradiction, that it is not 2-connected. Let c be a cut-vertex of G , and let $(A; B)$ be the corresponding decomposition of G .

Since every geodesic between vertices of A must lie in A , if there are $a, a' \in A$ with $d(a, a') = h$, the section graph of G defined by A is F -geodetic, a contradiction: therefore $d(a, a') < h$ for each $a, a' \in A$ and, similarly, $d(b, b') < h$ for each $b, b' \in B$.

Then there are $a' \in A, b' \in B$ such that $d(a', b') = h$: from now on we consider these vertices fixed.

Since G is not geodetic, let i be the *least* integer such that $F(i) \neq 1$. On a geodesic between a' and b' we choose $a \in A, b \in B$ with $d(a, b) = i$.

Every geodesic between a and b is the union of a geodesic from a to c and one from c to b . Since $d(a, c), d(c, b)$ are less than i , by our hypothesis on i both geodesics are uniquely determined. There is now a unique geodesic between a and b , contradicting the assumption $F(i) \neq 1$. This completes the proof.

We conclude with the following

EXAMPLE. The graph G in Fig. 2 is F -geodetic of diameter 4, with $F(2) = F(3) = 1$ and $F(4) = 2$; it is neither 2-connected nor minimally F -geodetic (namely the section graph of G defined by $V(G) \setminus \{1\}$ is F -geodetic).




FIGURE 2

ACKNOWLEDGMENT

The author thanks Professor P. V. Ceccherini for his careful reading of the manuscript in its original form.

REFERENCES

1. G. S. BLOOM, J. W. KENNEDY, AND L. V. QUINTAS, Distance degree regular graphs, in "The Theory and Applications of Graphs" (Proceedings, IV International Conf., Kalamazoo, Michigan), Lecture Notes in Mathematics, Springer-Verlag, New York, 1981.
2. J. BOSÁK, Geodetic graphs, in "Combinatorics" (Colloq. Kesthely, 1976), pp. 151-172, North-Holland, Amsterdam, 1978.
3. P. V. CECCHERINI, A q -analogous of the characterization of hypercubes as graphs, *J. Geom.* **22** (1984), 57-84.
4. A. GARDINER, Antipodal covering graphs, *J. Combin. Theory Ser. B* **16** (1974), 255-273.
5. D. B. HUGHES AND H. PIEPER, "Projective Planes," Springer-Verlag, New York, 1973.
6. W. M. KANTOR, Moore geometries and rank 3 groups having $\mu = 1$, *Quart. J. Math. Oxford Ser. (2)* **28** (1977), 309-328.
7. H. I. LEE, A note on geodetic graphs of diameter two and their relation to orthogonal latin squares, *J. Combin. Theory Ser. B* **22** (1977), 165-167.
8. M. MARCHI, Sottospazi lineari di spazi affini deboli, *Istit. Lombardo Accad. Sci. Lett. Rend. A* **110** (1976), 104-114.
9. O. ORE, "The Theory of Graphs," Amer. Math. Soc. College., Vol. 38, Amer. Math. Soc., Providence, R. I., 1962.
10. K. R. PARTHASARATHY AND N. SRINIVASAN, Geodetic blocks of diameter three, *Combinatorica* **4** Nos. 2-3 (1984), 197-206.
11. K. R. PARTHASARATHY AND N. SRINIVASAN, On geodetic blocks of diameter 2: Some structural properties, *Ars Combinatoria* **20** (1986), 49-60.
12. J. PLESNÍK, A construction of geodetic graphs based on pulling subgraphs homeomorphic to complete graphs, *J. Combin. Theory Ser. B* **36** (1984), 284-297.
13. J. G. STEMPLE, Geodetic graphs of diameter two, *J. Combin. Theory Ser. B* **17** (1974), 266-280.